

University of Wollongong

Research Online

Faculty of Science, Medicine and Health -
Papers: part A

Faculty of Science, Medicine and Health

30-1-2013

Sequential extraction procedures for the determination of phosphorus forms in sediment

Changyou Wang

Nanjing University of Information Science and Technology, changyou@uow.edu.au

Yong Zhang

Yantai Institute of Coastal Zone Research

Hongli Li

Nanjing University of Science and Technology

R John Morrison

University of Wollongong, johnm@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/smhpapers>



Part of the [Medicine and Health Sciences Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Wang, Changyou; Zhang, Yong; Li, Hongli; and Morrison, R John, "Sequential extraction procedures for the determination of phosphorus forms in sediment" (2013). *Faculty of Science, Medicine and Health - Papers: part A*. 876.

<https://ro.uow.edu.au/smhpapers/876>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Sequential extraction procedures for the determination of phosphorus forms in sediment

Abstract

This article summarizes the main extraction methods for sedimentary phosphorus (P) determination. With sequential chemical extractions, P is supposed to be selectively removed from different compounds in the sediments. Extraction schemes using strong acids and alkaline solutions have been tested on different sediments and found not to extract well-defined fractions. In addition, several systematic errors in these schemes have been detected. Thus, these schemes have been modified and simplified accordingly. The Standards Measurements and Testing Program of the European Commission (SMT) method is a popular modification of these extraction schemes, as it is simple to handle, allows laboratories to achieve reproducible results and could provide a useful tool for routine use by water managers. The SEDEX (sequential extraction method) method, another popular modification, is widely applied in biogeochemical research as it can separate authigenic carbonate fluorapatite from fluorapatite. Other chemical extractions using chelating compounds have attempted to extract P bound with iron and calcium in sediments without disturbing clay-bound or organic P, the purpose being to determine the algal-available non-apatite, apatite and organic fractions of sediment P. All extraction procedures still yield operationally defined fractions and cannot be used for identification of discrete P compounds. Future modifications of the extraction scheme should aim to achieve better extraction efficiency and selectivity, simple handling techniques and methods that can prevent the extracted P from being re-adsorbed onto Fe(OOH) and CaCO₃.

Keywords

extraction, procedures, sediment, phosphorus, sequential, determination, forms

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Wang, C., Zhang, Y., Li, H. & Morrison, R. John. (2013). Sequential extraction procedures for the determination of phosphorus forms in sediment. *Limnology*, 14 (2), 147-157.

Sequential extraction procedures for the determination of phosphorus forms in sediment

Changyou Wang^a Yong Zhang^b Hongli Li^a R. John Morrison^{c*}

a. School of Marine Sciences, Nanjing University of Information Science and Technology, Nanjing 210044, China

b. Key Laboratory of Coastal Zone Environmental Processes, Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences, Yantai 264003, China

c. School of Earth and Environmental Sciences, University of Wollongong, NSW 2522, Australia

Abstract: This paper summarizes the main extraction methods for sedimentary phosphorus (P) determination. With sequential chemical extractions, P is supposed to be selectively removed from different compounds in the sediments. The extraction schemes using strong acids and alkaline solutions have been tested on different sediments, and found not to extract well defined fractions. In addition, several systematic errors in these schemes have been detected. Then these schemes have been modified and simplified accordingly. The SMT (Standards Measurements and Testing Program of the European Commission) method is a popular modification of these extraction schemes, as it is simple to handle, suitable to allow laboratories to achieve reproducible results and could provide a useful tool for water managers on a routine basis. The SEDEX (sequential extraction method) method, another popular modification, is widely applied in the biogeochemical research as it is capable of separating authigenic carbonate fluorapatite (CFAP) from fluorapatite (FAP). Other chemical extractions using chelating compounds have attempted to extract P bound with iron and calcium in sediments without disturbing clay-bound or organic P, the purpose being to determine the algal-available non-apatite, apatite and organic fractions of sediment P. All extraction procedures still yield operationally defined fractions and cannot be used for identification of discrete P compounds. Future modifications of the extraction scheme should aim for better efficiency and selectivity in extraction, simple handling techniques and methods that could prevent the extracted P from being re-adsorbed onto Fe(OOH) and CaCO₃.

Key words: phosphorus, extraction, sediment, forms

1. Introduction

As a key component of fertilizer, phosphorus is crucial for the world's food supplies, and is intimately involved in both terrestrial and marine biogeochemical cycles (Blake et al., 2005; Jaisi and Blake, 2010). P is a potentially limiting nutrient that sustains primary productivity and has been recognized as a key factor responsible for eutrophication in lake, estuarine and some other waters (Schindler et al., 1973). The excessive supply of P (mainly from anthropogenic origin) to water systems, induces a degradation of their quality through the proliferation of algae, thus hindering various water uses. P in waters has two origins, i.e., external and internal. P from external origin comes from diffuse sources (natural, agricultural) or point sources (industrial and domestic

effluents). This nutrient is also released from the sediment, which acts as an internal source. Sediment can contribute phosphate to the overlying waters at levels comparable to the external source (Yuan et al., 2010). However, most of sediment P is in the particulate form, dissolved P being only a few per cent of total P (Ruban et al., 1999).

The recorded P mineral particulates include Apatite ($\text{Ca}_{10}(\text{F}, \text{OH})(\text{PO}_4)_6$), Brushite ($\text{CaHPO}_4 \cdot 2\text{H}_2\text{O}$), octocalciumphosphate ($\text{Ca}_3(\text{HPO}_4)_3 \cdot 3\text{H}_2\text{O}$), Anapaite ($\text{Ca}_3\text{Fe}(\text{PO}_4)_3 \cdot 4\text{H}_2\text{O}$), Strengite ($\text{FePO}_4 \cdot 2\text{H}_2\text{O}$), Vivianite ($\text{Fe}(\text{PO}_4)_2 \cdot 8\text{H}_2\text{O}$), Lipscombite ($\text{Fe}_3(\text{PO}_4)_2 \cdot \text{Fe}(\text{HO})_2$), Phosphoferrite ($(\text{Mn}, \text{Fe})_3(\text{PO}_4)_2 \cdot 3\text{H}_2\text{O}$), Ludlamite ($(\text{Fe}, \text{Mn}, \text{Mg})_3(\text{PO}_4)_2 \cdot 4\text{H}_2\text{O}$), Variscite ($\text{AlPO}_4 \cdot 2\text{H}_2\text{O}$), Wavellite ($\text{Al}_3(\text{PO}_4)_2(\text{OH}, \text{F})_3 \cdot 5\text{H}_2\text{O}$) (Pettersson et al., 1988). The P organic particulates, a complex fraction the exact nature of which is not precisely known, mainly consist of biogenic material. However, significant diagenetic reorganization of P occurs during burial, owing to changes in pH and redox potential. Not all the forms of particulate P are likely to be released, become bioavailable and, therefore, likely to increase eutrophication. The ability of sediment to store or release P has repercussions for the trophic state of waters, hence on their productivity. The study of the behaviour of P in the sediment is, therefore, a key factor which helps understanding the P cycle in rivers, lakes and other marine waters.

To understand the burial and diagenesis of P in sediments, it is necessary to identify, separate, and quantify the various solid-phase reservoirs of sedimentary P, which is difficult because of the low concentration and the fine-grained nature of most sediments. These two factors preclude use of most standard methods for direct identification of mineral phases, such as X-ray diffraction. Independent chemical analyses on phosphatic phases, such as isotopic or trace element signatures, clearly cannot be applied when discrete phases are not separable (Ruttenberg, 1992). The ^{31}P nuclear magnetic resonance (^{31}P NMR) can be used to assess the general composition of P in a variety of matrixes on both solid samples (solid state ^{31}P -NMR) and on samples in solution (high resolution ^{31}P -NMR). Though it is possible to learn something about solid-phase P in sediment using solid state ^{31}P -NMR, this is seldom used due to the inherently low sensitivity of NMR and low natural abundance of P in sediment. The high-resolution NMR is the better method because it allows pre-concentration of the sample before analysis, but high resolution ^{31}P -NMR also requires extraction of P from the original solid matrix (Ahlgren et al., 2006, 2007). As a result, it is necessary to use fractionation of inorganic P to identify the amount in each discrete chemical form.

The various forms in which P occurs in sediments have received considerable attention during recent decades and many extraction schemes have been developed. The methods for the chemical extraction of different forms of sediment P originated from soil science. Sequential extractions were designed to quantify step by step discrete chemical or mineralogical compounds. Extraction methods not only provide information on the potential mobility of P species, they also allow the assessment of the origin of the sediment P. The best example of these early extraction schemes is the one presented by Chang and Jackson in 1957. The Chang and Jackson procedure was further developed by Williams et al. (Williams et al., 1967; 1971 and 1976), Hieltjes et al. (Hieltjes and Lijklema, 1980), Psenner et al. (Psenner and Pucsko, 1988), Ruttenberg et al. (1992), Golterman et al. (1982, 1995, 1996, 1998). These authors pointed out the shortcomings of the former procedure and suggested their own modifications. However, to date, no standardised method is available for the determination of different forms of P in sediments. These P forms determined using sequential extraction procedures are operationally defined and are related to specific reagents and procedures, such as authigenic carbonate fluorapatite (CFAP) separated from detrital apatite by acetate buffer,

i.e., results are interpreted to be related to a specific phase of sediment (although, *sensu stricto*, it is related solely to a chemical procedure). It has been recognised that the different P fractions are of different significance in terms of ecological meaning, and this knowledge is also updated gradually with the use of new technology or methods: for example, the belief that P extracted by NaOH reflects the P most available to algae (Williams et al., 1980), was questioned by Golterman et al. (1996), who stated that ethylene diammine tetraacetic acid (EDTA) as a P extractant (instead of NaOH) gave more information on the quantity of P available for algae.

This paper summarizes the main extraction methods for sedimentary P determination, including Williams method (Williams et al., 1967; 1971 and 1976), Hieltjes–Lijklema method (Hieltjes and Lijklema, 1980), Golterman method (Golterman et al., 1982, 1995 and 1996), sequential extraction (SEDEX) method (Ruttenberg, 1992), standards measurements and testing method (SMT) (Ruban et al., 1999 and 2001), Penn method (Penn et al., 1995 and 1997), Jensen method (Jensen and Thamdrup, 1993; Jensen, 1998), discusses the ecological significance of the fractional composition, comments on the inherent problems and applicability of chemical fractionation, and makes a practical recommendation.

2. Development of sequential extraction procedures of P forms in sediments

Sequential extraction methods take advantage of the fact that different solid phases show dissimilar reactivity toward different solutions. Sediment is extracted with a series of extractants, each chosen to selectively dissolve a single phase or group of phases of similar chemical characteristics. The sequence is designed so that the most reactive phases are removed first and the severity of the extractants increases with each step. Sequential extraction schemes were first developed for soils and then extended to sediments. Characterization of the different forms of inorganic phosphorus in soil by sequential extraction with acid and alkaline reagents was initiated by Dean in 1938, and later modified by Chang and Jackson in 1957. Most procedures used in sediments are modifications or improvements of the sequence, based on the extraction using acid and alkaline reagents, originally suggested by Chang and Jackson (Chang and Jackson, 1957; Tiyaopongpattana et al., 2004). There are no radical differences in extractants between soil and sediment extraction scheme except for the specific method for sediment to distinguish authigenic apatite from detrital apatite and pretreatment approaches to treat soils and sediments prior to extraction (Williams et al., 1971; Hieltjes and Lijklema, 1980; Ruttenberg, 1992; Condrón and Newman, 2011). Most operationally defined schemes available address inorganic forms of phosphate. The main inorganic forms of P are the fraction adsorbed by exchange sites and referred to as loosely bound, labile or exchangeable P, the fraction associated with Al, Fe and Mn oxyhydroxides and the fraction in Ca-bound compounds generally referred to as apatite P or Ca-bound (Ruban et al., 1999b). Organic P is a complex fraction the exact nature of which is not precisely known and is normally taken into account as a whole. The steps involved in each method are summarized in Table 1.

2.1 Williams method

The Chang and Jackson procedure using acid and alkaline reagents was modified and applied to the sediment by Williams and co-workers. The shortcomings of the original procedure were pointed out, which included the extraction of iron-bound P plus aluminum-bound P with ammonium

fluoride, as well as the re-adsorption of phosphate due to formation of calcium difluoride, especially in calcium-rich sediments. They therefore suggested several modifications (Williams et al., 1967; 1971 and 1976) as indicated in Table 1.

Williams et al. (1976, 1980) used an extraction method with CDB (tri-sodium citrate, sodium dithionite and sodium bicarbonate) followed by NaOH. A subsequent extraction with HCl is supposed to extract Ca-P. The shortcomings of these two versions have already been outlined including re-adsorption of phosphate onto carbonate that could occur in the NaOH extraction, especially in calcareous sediments (Hieltjes and Lijklema, 1980). It was also shown that a substantial amount of calcium-bound P was solubilized by the citrate dithionite bicarbonate (CDB) reagent of Williams (1971, 1976), for which the strongly chelating citrate ion was responsible. The CDB extraction was omitted by Burrus et al. (1990), but the name of the modified procedure was not changed and usually listed as a later version of the Williams scheme (Golterman, 1996; Ruban et al., 1999b; Hoyle, 2004). The reason for this change was the belief that P extracted by NaOH reflects the P available to algae. The idea of identifying discrete P compounds was gradually abandoned and sediment P content was finally separated into non-apatite P, apatite P and organic P (OP). The non-apatite P was considered as bioavailable, apatite P non-available and OP partly available. OP was considered to be completely extracted by hydrochloric acid and sodium hydroxide, the latter in two consecutive steps. Total P (TP) and molybdate reactive P were measured in the pooled extracts and OP calculated from the difference (Williams et al., 1976). The extraction scheme of Williams as modified by Burrus et al. (1990) is not completely sequential, involving two independent procedures.

Table 1. Extraction schemes for determining the fractional composition of sediment phosphorus.

Procedure	Extraction	Proposed fraction	Advantages	Shortcomings
Chang and Jackson (1957)	a. NH_4Cl 1 M	labile P	identifying discrete P compounds	extraction of Fe-P by NH_4F resorption by CaF_2 precipitation of phosphate with iron
	b. NH_4F 0.5 M pH 8.2	Al-bound P		
	c. NaOH 0.1 M	Fe-bound P		
	d. HCl 0.5 M	Ca-bound P		
	e. CDB	reductant-soluble P		
	f. NaOH	refractory P		
Williams (1976)	a. CDB 0.22/0.1 M	non-apatite P	Simple, practical	resorption by carbonates in calcareous sediments Ca-P and org-P released by CDB
	b. NaOH 1 M	apatite P		
	c. HCl 0.5 M	organic P		
Williams (Burrus et al., 1990)	a. NaOH 1 M	non-apatite P	Simple, practical	partial resorption of P extracted by NaOH on CaCO_3
	b. HCl 1+3.5 M	apatite P		
	c. HCl 3.5 M+calcination	total P		
	d. HCl 1 M +calcination	organic P		
Hieltjes and Lijklema (1980)	a. NH_4Cl 1 M pH 7	labile P	Simple, practical	dissolution of small amounts of Fe-P and Al-P by NH_4Cl ; hydrolysis of organic P; no relation with bioavailability
	b. NaOH 0.1 M	Fe- and Al-bound P		
	c. HCl 0.5 M	Ca-bound P		
Golterman (1996)	a. H_2O	labile P bioavailable	giving more information on the quantity of P available	not practical; EDTA interfere with P determination; complicated solution preparation; in some sediments,
	b. Ca-EDTA 0.05 M + dithionite	iron-P bioavailable		
		Ca-P non-available		

	c. Na ₂ -EDTA 0.1 M	acid soluble organic P	for algae	extraction must be repeated
	e. H ₂ SO ₄ 0.25 M	bioavailable		
	f. NaOH 2 M	reductant Organic P		
		non-available		
SEDEX (Ruttenberg, 1992)	a. MgCl ₂ 1 M	loosely sorbed P	separating authigenic	very long; not practical; partial OP
	b. CDB 0.3/1 M	ferric Fe-bound P	CFAP from detrital	released by CD
	c. Acetate buffer 1 M	CFAP+ biogenic apatite+ CaCO ₃ -P	FAP	
	e. HCl 1 M	detrital apatite		
	f. calcination+HCl 1M	organic P		
SMT (Ruban et al., 2001)	a. NaOH 1 M (extract+3.5 M HCl)	NaOH-P	Simple, practical	partial resorption of P extracted by NaOH on CaCO ₃
	b. 1 M HCl	HCl-P		
	c. HCl 1 M	IP		
	d. HCl 1 M +calcination	OP		
	e. HCl 3.5 M +calcination	Conc. HCl-P		
Penn (1995)	a. NH ₄ Cl 1 M (3×)	loosely bound-P	dissolution of small	hydrolysis of organic P; no relation
		loosely sorbed and CaCO ₃ associated-P	amounts of Fe-P and Al-P by NH ₄ Cl was	with bioavailability
	b. NaOH 0.1 M (split)	Fe and Al bound-P	avoided	
	c. NaOH 0.1 M (split)	extractable biogenic-P		
	d. HCl 0.5 M	calcium mineral-P		
	e. Persulfate digestion	refractory organic-P		
Psenner (1988)	a. NH ₄ Cl 1 M	soluble P	identifying discrete P	resorption by carbonates in
	b. DB 0.11 M 40 °C	reductant-soluble P	compounds	calcareous sediments; hydrolysis of organic P
		Fe- and Al-bound P		
	c. NaOH 1 M	refractory OP		
	d. NaOH 1M	Ca-bound P		
	e. HCl 0.5 M			
Jensen (1998)	a. MgCl ₂ 1 M	loosely adsorbed IP and leachable OP	separating authigenic	very long; not practical; resorption
			CFAP from detrital	by carbonates in calcareous
	b. DB 0.11 M	iron-bound IP	FAP	sediments
		adsorbed IP and leachable OP		
	c. NaOH 0.1 M	authigenic apatite, CaCO ₃ -bound IP and leachable OP		
	d. Acetate buffer 1M	detrital apatite		
		refractory OP		
	e. HCl 0.5 M			
	f. ignition+HCl 1 M boiling			

2.2 Hieltjes–Lijklema method

The need for chemical characterisation was stressed in the Hieltjes and Lijklema method in order to improve comprehension of the P exchange processes between sediment and water and the effects

of environmental conditions such as pH, redox potential and ionic strength on these processes (Hieltjes and Lijklema, 1980).

The Hieltjes–Lijklema method uses NaOH to solubilise Fe and HCl to dissolve Ca and the NaOH extraction is preceded by NH₄Cl extraction. In this method, residual P is given by the difference between the total P content and the sum of reactive (inorganic) fractions and mainly consists of organic P. Although the same reagents are used in the Williams and Hieltjes–Lijklema methods, their concentrations differ; they are higher in the Williams method. The Williams method yields OP, which is not the case with the Hieltjes–Lijklema method (OP can be calculated as the difference between TP and IP). The Hieltjes–Lijklema method yields soluble labile P, whereas this phase is not determined in the Williams scheme. Also, the Hieltjes–Lijklema method is a sequential extraction scheme, whereas in the Williams protocol the steps are independent, not sequential.

It is considered an advantage of the Hieltjes and Lijklema method when compared with the Williams method, that only one sediment sample is needed. However, the concentration of the NaOH extractant and the influence of the extraction duration have not been taken into account sufficiently in this method. Repeating the treatment with NaOH, Golterman (1996) found that a second extraction may extract another~30% of this P pool, while increasing the NaOH concentration again extracted more orthophosphate. It is obvious that what is extracted with the NaOH is certainly not only Fe-bound P (Golterman, 1996). The other shortcomings of this method include that dissolution of small amounts of Fe-P and Al-P by NH₄Cl was possible in the Hieltjes–Lijklema scheme. Furthermore, though the back precipitation onto CaCO₃ in the NaOH extraction had been taken into account, the improvement in removing the calcium carbonate by ammonium chloride extraction was inefficient, especially in highly calcareous sediments. This procedure does not extract specific compounds and gives, therefore, no information on the quantity of P available for algae if CaCO₃ and/or org-P were present in reasonable amounts (Golterman, 1996).

2.3 Golterman method

Golterman et al. (1982, 1995 and 1996) have argued that strong acids and alkaline solutions should be avoided in P extractions schemes since they are too aggressive and do not deliver well defined fractions. By using a chelating compound, Fe- and Al-bound P would be extracted without disturbing clay-bound or organic P.

Several modifications had been suggested by Golterman et al. (1982, 1995 and 1996). At first, repeated extractions with 0.01 M NaNTA (sodium nitrilo triacetate) solution were suggested, but later sequential dissolution of Fe- and Ca-bound P with 0.05 M CaNTA (calcium nitrilo triacetate) and 0.05 M NaNTA, respectively, has been proposed (Golterman, 1982). NTA (Nitrilo triacetic acid or its salts) is replaced subsequently by EDTA. The EDTA fractionation of sediment-bound inorganic phosphate was improved and made more effective. The Fe-bound phosphate in a recent version was extracted with Ca-EDTA plus dithionite and followed by an extraction of Ca-bound phosphate by Na₂-EDTA at pH =4.5 (Golterman, 1996). The method using EDTA as P extractant instead of NaOH, is different from the Williams and Hieltjes–Lijklema methods, and the Fe-bound phosphate extracted with Ca-EDTA gives more information on the quantity of P available for algae than the P extracted by NaOH. NaOH is too aggressive and does not deliver well defined fractions (Golterman, 1996), as the pH during the extraction is very different from the pH of the sediment, causing changes in the original P composition. Fabre et al. (1996) have shown that P extracted by

NaOH is not equal to the P available for algae. Dorich et al. (1985) demonstrated a strong correlation between algal available P and P extracted by NaOH, but also that the NaOH extraction largely overestimated the available P. It seems that this overestimation is caused by the hydrolysis of non-available OP

The Golterman method is not without pitfalls. EDTA solutions have the disadvantage of interfering with the orthophosphate and Fe determinations. However, Golterman insisted that for the orthophosphate determination 2 ml 0.1 M of EDTA could be used without interference (Golterman, 1996). Another disadvantage of the NTA/EDTA extractions is that they must be repeated, as one single extraction can never give complete recovery.

2.4 SEDEX

The Ruttenberg method, i.e., SEDEX scheme, initially developed for marine sediments, is a five-step procedure which is capable of separating authigenic CFAP from detrital apatite of igneous or metamorphic origin (FAP) and is widely applied in biogeochemical research.

The SEDEX method uses MgCl_2 to extract loosely sorbed or exchangeable P, CDB to solubilise ferric Fe-bound P, acetate buffer to dissolve authigenic carbonate fluoroapatite and biogenic apatite and CaCO_3 associated P, HCl to dissolve detrital apatite and a second HCl extraction after calcination to measure organic P. The SEDEX method is the first method that has been devised to chemically separate two of the main categories of authigenic phosphate phases called upon most often as sedimentary sinks for diagenetically mobilized P: ferric oxyhydroxide-associated P (Fe-P) and authigenic CFAP. It also offers a means for separating authigenic CFAP from FAP, which is the main advantage of the SEDEX method. The importance of this distinction is that CFAP represents an oceanic sink for reactive P, whereas detrital apatite does not. Coupled with pore-water and other solid-phase data, the SEDEX method is capable of revealing otherwise inaccessible information on the chemical nature of sedimentary P. However, although Ruttenberg (1992) also stated the problem of analytical artifacts resulting from redistribution of P onto residual solid surfaces during extraction had also been resolved by washing the sediment residue with MgCl_2 between extraction steps, no data was found to support this claim. The five-step SEDEX was streamlined to a four-step procedure by Anderson and Delaney (2000). The major modifications were the combination of extraction of adsorbed and oxyhydroxide-associated pools into a single step and the use of a lower dithionite concentration, resulting in decreased analytical background and working days required. In addition, automated spectrophotometric flow injection analysis was used to determine P concentrations, rather than traditional spectrophotometric techniques.

The underestimates for OP values due to SEDEX pre-extraction of organic P in the prior steps had been discussed by Ruttenberg (1992). Pre-extracted organic P will not affect the accuracy of the measured quantities of P in steps before calcination if it remains unhydrolyzed. At least a portion of organic P extracted by HCl remains unhydrolyzed (Ruttenberg 1990). Golterman (1996) had shown that the SEDEX extraction gave results not different from those with Ca-EDTA or Ca-NTA, but extracted less CaCO_3 -P. This difference cannot be explained with certainty, but it seems likely that the CDB extracted some CaCO_3 associated P. The CDB reagent was found to be not sufficiently effective in separating Fe-P from CaCO_3 -P, especially for calcareous sediments, presumably because citrate is a strong complexing agent for both Fe and Ca (Hieltjes and Lijklema, 1980). Furthermore, citrate has also been found to interfere with spectrophotometric determination of phosphate. For these reasons, Zhang et al (2004) made an important modification to the SEDEX

scheme by omitting citrate from the CDB step when they studied sedimentary phosphorus in Florida Bay. The fact that iron (oxy)hydroxides are not completely extracted makes a re-adsorption onto iron (oxy)hydroxides likely. In fact, some forms of apatite behave differently (Barbanti et al., 1994). The SEDEX method must be tested to see whether repetition of an extraction (especially that with acetate buffer) would yield some extra P (Golterman, 1996).

2.5 SMT

Although the various forms of P in sediments have received considerable attention in recent decades and many extraction schemes have been developed, there is no standardized method for P extraction. However, the comparability of data is only possible on the basis of standardized procedures that require to be collaboratively tested and agreed by group(s) of researchers. In order to improve this situation, the European Commission through the Standards, Measurements and Testing Program launched a collaborative project to design a harmonized sequential extraction scheme.

This scheme, the so-called SMT extraction scheme, was a modified version of the Williams protocol, which involved three independent procedures, yielding NaOH-P, HCl-P, IP, OP and TP. The SMT protocol had been further tested on three sediments (calcareous, siliceous, organic rich) and the results reported were considered satisfactory (Ruban et al., 2001). The SMT and the Golterman schemes were compared for a sediment of the Bort-Les-Orgues Reservoir, France. The SMT method appeared to be more satisfactory than the Golterman method, as the Golterman method was mainly designed for calcareous sediment and the Bort-Les-Orgues sediments are siliceous (Ruban et al., 1999a).

The SMT method is simple to handle, suitable for laboratories to achieve reproducible results and could provide a useful tool for water managers on a routine basis, especially at a time when quality assurance and data comparability are of paramount importance in laboratory analysis. Moreover, the method could help in calculating the releasable P stock in lake sediment, and hence provide information on the lake recovery delay (Ruban et al., 2001). However, it should be noted that the SMT protocol was not better with regard to the evaluation of different organic P fractions, information on bioavailable fractions, and distinction between different apatite forms. It would appear to be most useful as a tool for management.

2.6 Other modifications of extraction procedures

2.6.1 Penn's modification of the Hieltjes and Lijklema method

Penn et al. (1995, 1997) proposed a sequential chemical extraction method to quantify the P fractions, which had been applied to the sediment in Onondaga Lake, a calcareous, eutrophic lake in Syracuse, New York, U.S.A.

This method was a modification of original procedure of Hieltjes and Lijklema (1980) that called for two consecutive extractions with NH_4Cl followed by single extractions with NaOH and HCl. In the Penn et al. (1995) scheme, a third NH_4Cl extraction was added for highly calcareous sediments, and NaOH extracts were split for reactive P (NaOH-RP) and non-reactive P (NaOH-NRP) determinations (NaOH-NRP is determined as the difference between TP and NaOH-RP). NaOH-NRP includes easily hydrolysable organic P and polyphosphates (Hupfer et al., 1995) and is thought to be an indicator of biologically available P.

This technique does not appropriately differentiate the Fe- and Al-P and CaCO_3 -P fractions on oxidized samples due to adsorption of solubilized P to iron (oxy)hydroxides during the NH_4Cl

extraction. Here the two fractions are combined and termed loosely bound-P. This analytical artifact does not influence estimates of labile-P which are based on the sum of Fe and Al-P, CaCO_3 -P, and extractable biogenic-P. The shortcoming in the Hieltjes and Lijklema (1980) scheme, that dissolution of small amounts of Fe-P and Al-P by NH_4Cl was avoided. However, the concentration of the NaOH and the influence of the extraction time had not been taken into account in this method either.

2.6.2 Psenner's modification of Chang and Jackson method

The fractionation of P scheme recommended by Psenner (Pettersson et al., 1988) agrees with the original ideas of Chang and Jackson (1957) about finding chemical or mineralogical definitions of the P forms in sediments, although the authors are aware of the difficulties and restrictions involved. In the later versions of the Psenner et al (1988) scheme, a 1 M NH_4Cl extraction was used, instead of water for extraction of loosely adsorbed P in the first step (Psenner and Pucsko 1988). Kerr et al. (2010) proposed a modification of Psenner et al (1988) scheme and applied this to a subtropical river sediment in Australia. A subsample of extract was digested with 1 M NaOH using the persulfate digestion before analysis, leading to the determination of non-reactive NaOH-extractable P (NaOH-NRP), which is thought to represent labile organic and bacterial P.

This extraction scheme was originally developed for siliceous sediments, and re-adsorption of P by carbonates was unavoidable in calcareous sediments (Psenner and Pucsko 1988; Jensen and Thamdrup 1993).

2.6.3 Jensen's modification of Psenner method

A sequential six-step extraction scheme for P pools was developed by Jensen and co-workers (Jensen and Thamdrup, 1993; Jensen, 1998) for use in marine sediments which permitted an operational discrimination between seven different P pools (see Table 1).

The extraction scheme was a modification of the scheme proposed for freshwater sediment by Psenner et al. (1988). NH_4Cl in the Psenner (1988) scheme was found to extract more Ca and soluble reactive P. This indicates dissolution of CaCO_3 -P. The major modifications were the use of 1 M MgCl_2 instead of 1 M NH_4Cl for extraction of loosely adsorbed P in the first step and the inclusion of an additional acetate buffer extraction after extraction with NaOH and before the HCl extraction. Both modifications were adapted from Ruttenberg (1992). Moreover, a NaCl wash was added after each extraction step in order to minimise the re-adsorption of phosphate. With this modified extraction procedure, the loosely adsorbed IP, along with some OP, was intended to be extracted in the first step. The dithionite reduces oxidized Fe and Mn species and dissolves both the metals and the associated IP (Fe-P). The BD-reagent was found to be very specific for Fe-P and BD-extractable P was by far the most exchangeable P pool even deep in the sediment where the pool size was small (Jensen and Thamdrup, 1993). The NaOH extraction was intended to extract a large proportion of OP and remove IP adsorbed onto surfaces other than those of reducible metal oxyhydroxides (e.g. clay minerals, gibbsite). IP associated with humic acids is also extracted in this step. The acetate buffer dissolves the biogenic carbonates and associated CFA. Detrital apatite will dissolve in 0.5 M HCl. Combustion followed by hot HCl extraction dissolves and hydrolyzes the residual OP compounds (Jensen, 1998).

This modified extraction scheme was also originally developed for siliceous sediments, so P re-adsorption by carbonates was unavoidable in calcareous sediments (Psenner and Pucsko 1988;

3. Ecological interpretation of operationally defined P fractions

Generally, bioavailable or mobile P in the sediment can be defined as the sum of immediately available P and potential P that can be transformed into an available form by naturally occurring physical, chemical and biological processes (Wang et al., 2009). The exchangeable or loosely sorbed phosphate is available for algal growth according to bioassays. The iron-bound phosphate is potentially bioavailable, depending primarily on sediment redox intensity (Katsaounos et al., 2007). The organic P could become bioavailable by microbial remineralization.

Since P has been identified to be the primary limiting factor for algal growth in many natural waters several decades ago (Ruban et al., 1999a), a considerable number of researchers have focused their work on sedimentary P speciation and on water-sediment interactions (Lukkari et al., 2007). One of the main aims of these studies was to assess the bioavailability of sedimentary P within the range of natural physical and chemical characteristics in water. In other words, it is important to know whether certain forms of P can be released from the sediment, under certain conditions, into the water column for subsequent algal development. Among the various P-containing minerals, calcium salts (mainly apatites, generally coprecipitated with calcium carbonate) are probably the most inert in the slightly alkaline, poorly complexing, normal aquatic conditions. Dissolution of these minerals implies a sharp pH decrease. Iron-bound phosphate is both pH and redox dependent since ferric hydroxide precipitates are involved in its formation. Under anoxic conditions, generated by intense organic matter mineralisation, Fe(III) is reduced into Fe(II), which dissolves in the water phase along with associated phosphate. Bioavailability of organic P requires bacterial mineralization into phosphate (Anderson et al., 2010).

By measuring uptake of sediment P by algae in culture and extraction by NaOH and nitrilotriacetic acid solutions and by H-resin, Williams et al. (1980) tested several extraction schemes on eroding bluff materials and lake and fluvial sediments from the Lake Ontario drainage basin, aiming to determine which of the fractions of sediment P was most available to algae and to relate these fractions to the amounts of P removed by various chemical extractants which had been proposed for measuring “available P”. They found that P uptake by the algae was related to the amount of non-apatite inorganic P in the sediments. Apatite P was not consumed. The non-apatite inorganic P fraction was highly correlated with the amounts of inorganic P extracted by three techniques for estimating “available P” and cell uptake equaled NaOH-extractable inorganic P in several instances. Organic P in the sediments was not utilized by the algae (Williams et al., 1980).

Interest in organic P was, however, renewed with the use of ^{31}P NMR spectroscopy since the 1980s. The chemical forms of organic P were identified, such as orthophosphate monoesters, diesters, or phosphonates. Microbial P lipids and orthophosphate monoesters decompose rapidly in sediments. Some forms of organic P, such as teichoic acid P, are barely decomposed. The degradation of organic P compounds in the sediment is substantial and may be an important source of bioavailable P to the water column (Ahlgren et al., 2006, 2007). Owing to the lengthy nature of algal assays, an alternative technique using iron oxide-impregnated filter paper strips (Pi-strip) was developed in 1980s. In this, iron oxyhydroxide precipitate on paper strips reacts with phosphate ions in solution forming water-insoluble iron phosphates on the paper which are subsequently removed in dilute acidic solution. The strips therefore act as P sinks, approximating to P removal from water

by algae (Sharpley, 1993a, b). Sharpley (1993a,b) compared the Bioavailable P content determined by the Pi-strip method to the growth of P-starved algae and found that the two methods were strongly correlated and that in turn, P determined by NaOH extraction produced a 1:1 relationship with Pi-strip P.

However, it has been proposed that not only is the NaOH extractant concentration important, but that the duration of the extraction also influences the quantity extracted. It has also been shown that when NaOH-P largely overestimated the P available for algae, the NaOH extract was certainly not removing Fe-bound P only (Dorich et al., 1985; Fabre et al., 1996). It seems very probably that this overestimation is caused by the hydrolysis of non-available OP. If, as in the Williams method, NaOH is used as P-extractant, the method is not different from Hieltjes–Lijklema method, which is essentially the old Chang & Jackson method with a preliminary washing with NH_4Cl at $\text{pH} = 7$ the function of which is rather uncertain. The presence, in all sediments, of different forms of OP which are partly hydrolysed by NaOH and HCl, should be taken into account by geochemists, as these compounds behave biologically different from the iron- and calcium-bound phosphates.

By comparing the BD extractable P-pool with both the pool of iron in the BD-fraction and the pool of oxidized, amorphous or poorly crystalline iron oxyhydroxides, highly significant correlations ($p < 0.001$) were observed; Jensen and Thamdrup (1993) concluded that the BD-reagent was very specific for iron-bound P. Further evidence for this came from two experiments: 1) enhanced BD treatment did not result in additional phosphate extraction; and, 2) by sequential extraction of P pools in pure cultures of diatoms and cyanobacteria no phosphate was recovered in the BD-fraction. Further, an isotopic exchange experiment with ^{32}P revealed that BD-extractable P was by far the most exchangeable P-pool even deep in the sediment where the pool size was small (Jensen and Thamdrup, 1993).

Golterman et al. (1996) stated that EDTA as P extractant instead of NaOH, which was different from the Williams and Hieltjes–Lijklema methods, gave more information on the quantity of P available for algae, and that those authors who did not obtain a good extraction with Ca-NTA/EDTA was due to the use of a Ca-NTA/EDTA extractant for a very short time only and without a reducing agent. Ruban et al. (2001) also believed inorganic P extracted by the Ca-EDTA gave more information on bioavailable fractions.

4. The main problems with sequential extraction methods for sediment P and a practical recommendation

4.1 Extractant

Strong acids and alkaline solutions should be avoided in P extractions schemes since they are too aggressive and do not deliver well defined fractions. By using a chelating compound, such as EDTA, iron- and aluminum-bound P would be extracted without disturbing clay-bound or organic P; however, EDTA solutions have the disadvantage of interfering with the orthophosphate and Fe determinations. There is an urgent need to identify a “New” extractant with better efficiency and selectivity. More attention should also be paid to pH as it strongly influences P speciation in sediment. The extractants should be different for the different sediments (calcareous, siliceous, organic rich) and be standardized to achieve better efficiency and selectivity, but not the extraction conditions such as duration of the extraction and the concentrations of the extractants, as these depend strongly on the nature of the sediments, the concentrations of the different P fractions and on

the solid:solution ratio.

4.2 Redistribution

The second problem is the analytical artifacts resulting from redistribution of P onto residual solid surfaces during extraction. Ruttenberg (1992) stated that the secondary adsorption onto residual solid surfaces during extraction could be successfully reversed by washing the sediment residue 2 times with MgCl_2 between extraction steps. This method is time consuming and the many steps are a source of cumulative error in spite of the detailed protocol. Any new methods should be simple to carry out, and able to prevent completely re-adsorption of extracted P onto $\text{Fe}(\text{OOH})$ and CaCO_3 .

4.3 Pretreatment of the sediment

It is obvious that in many laboratories, the environmental conditions under which samples for P analysis are stored are different from those in the natural sediments, especially the content of water and oxygen. The OP and Fe-P would change in quantity with the changes in pH and redox potential. P produced by organic matter decomposition would be redistributed by adsorption onto iron oxyhydroxide in the oxic atmosphere. Golterman (1996) stated that fresh sediments should be used, as freezing or drying changed the fractionation results. The disadvantage for a methodological study is that not all sediments are intercomparable in water and dissolved oxygen. Monbet et al. (2007) had tried to improve the pretreatment. They made sediment cores sectioned in a homemade glove box filled with argon. Each slice was immediately centrifuged under argon. After the removal of pore water, the sediment samples were freeze-dried. All sediment slicing and pore-water extractions were directly performed on site immediately after core collection in order to minimise disturbance of the core. By this procedure, the impact of oxygen on sediment P could be diminished; however, the fractionation results could still be changed in the freeze-drying step.

4.4 Recommendation

There is still not a widely acceptable and standardized method for the determination of P speciation due to its variety and changeability in sediments. With the many types of naturally occurring P classified via only a few extraction steps, chemical fractionation methods cannot be more than an approximate approach. However, determination of the chemical character of sediment P still allows a general estimate of the portion of sediment P that, under certain environmental conditions, can become available for primary production in aquatic environments. At the same time, sequential extraction provides useful information on the proportions and forms of P that do not participate in the nutrient cycle but are buried with deposited material.

As to the different sediments, a suitable extraction scheme should be selected in order to acquire useful information. The Williams (1976, 1990) method was originally developed by modifying the Chang and Jackson soil P extraction scheme and applied to lake sediments. This method gives better results for siliceous sediments and the non-apatite P extracted by NaOH can be considered as bioavailable though it often leads to overestimates (Williams et al., 1976; Burrus et al., 1990). This method is not suitable to be used in calcareous sediments because of re-adsorption by carbonates (Hieltjes and Lijklema, 1980; Golterman, 1996; Pettersson et al., 1988).

As the NaOH extraction is preceded by a NH_4Cl extraction, the Hieltjes and Lijklema (1980) method could be applied not only to siliceous sediments but also to calcareous lake sediments

(Hieltjes and Lijklema, 1980). However, the extracted P was not related with bioavailability if any substantial amount of CaCO_3 and/or org-P was present (Golterman, 1996). Moreover, the capacity of CaCO_3 removal by the NH_4Cl extraction was limited, especially in highly calcareous sediments. This method can be a good choice in siliceous sediments or calcium-poor sediments when the information on chemical characterisation of the extracted P and the P exchange processes between sediment and water is needed (Hieltjes and Lijklema, 1980).

When the data quality of bioavailable P and OP are of paramount importance, the Golterman (1996) method is the best option for calcareous sediments (Ruban et al., 1999a), in which Fe-bound phosphate extracted with Ca-EDTA gives more information on the quantity of P available for algae and no pre-solubilization of OP resulted in underestimates for OP values (Golterman, 1996).

The SEDEX (1992) method was originally developed for use with marine sediments (Ruttenberg 1992) and is now also used for sequential extraction of P from freshwater sediments (Ruban et al. 1999b; Simon et al. 2009). This method is the best option for the biogeochemical research as it is capable of separating CFAP from FAP, which is an important distinction as authigenic CFAP represents an oceanic sink for reactive P and detrital FAP does not.

The SMT (2001) method is simple to operate, suitable for laboratories to achieve reproducible results and therefore a popular extraction scheme in waters management (Ruban et al., 2001). It gives no information, however, on bioavailable fractions, especially for highly calcareous lake sediments, due to re-adsorption of P extracted by NaOH on CaCO_3 , and also no distinction between different apatite forms. This method could be a best option for siliceous sediments when a useful tool for water managers on a routine basis is needed.

The modifications to extraction procedures have improved the original procedures. For example, the Penn et al. (1995) scheme, a modification of the Hieltjes and Lijklema (1980) method, could be used in highly calcareous sediments by adding a third NH_4Cl extraction before extractions with NaOH and HCl. Jensen's modification (1993), mainly by an additional acetate buffer extraction after extraction with NaOH and before the HCl extraction in the Psenner (1988) method, can be used for the biogeochemical research as it is capable of separating CFAP from FAP, but would be most suitable for siliceous sediments as re-adsorption by carbonates is unavoidable in calcareous sediments (Psenner and Pucsko 1988; Jensen and Thamdrup 1993). Endeavors at modification of sequential extraction procedures for determination of P are ongoing due to the necessity in sedimentary P research and we look forward to the development of even better techniques.

Acknowledgements

This study was supported by the National Natural Science Foundation of China (Project No. 41006040) and by the government Scholarship of Jiangsu Province, China for study abroad and by the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD).

References

- Anderson L D and Delaney M L. Sequential extraction and analysis of phosphorus in marine sediments: Streamlining of the SEDEX procedure. *Limnol. Oceanogr.*, 2000, 45(2): 509–515.
- Anderson L D, Faul K L, Paytan A. Phosphorus associations in aerosols: What can they tell us about P bioavailability? *Mar. Chem.*, 2010, 120: 44–56.

- Ahlgren J, Brabandere H D, Reitzel K, Rydin E, Gogoll A and Waldebäck M. Sediment Phosphorus Extractants for Phosphorus-31 Nuclear Magnetic Resonance Analysis: A Quantitative Evaluation. *J. Environ. Qual.* 2007, 36: 892-898.
- Ahlgren J, Reitzel K, Tranvik L, Gogoll A and Rydin E. Degradation of organic phosphorus compounds in anoxic Baltic Sea sediments: A ^{31}P nuclear magnetic resonance study. *Limnol. Oceanogr.*, 2006, 51(5):2341-2348.
- Aminot A. and Andrieux F. Concept and determination of exchangeable phosphate in aquatic sediments. *Water Res.*, 1996, 30 (11): 2805-2811.
- Barbanti A, Bergamini M C, Frascari F, Miserocchi S, Roso G. Aspects of Sedimentary Phosphorus Chemical Fractionation. *J. Environ. Qual.*, 1994, 23: 1093-1102.
- Blake R E, O'Neil J R and Surkov A. Biogeochemical cycling of phosphorus: insights from oxygen isotope effects of phosphoenzymes. *Am. J. Sci.* 2005, 305: 596-620.
- Burrus D, Thomas R L, Dominik B, Vernet J-P, Dominik J. Characteristics of suspended sediment in the upper rhone river, switzerland, including the particulate forms of phosphorus. *Hydrol. Process.*, 1990, 4 (1): 85-98.
- Chang S C, Jackson M L. Fractionation of soil phosphorus. *Soil Sci.*, 1957, 84: 133-144.
- Condrón L M and Newman S. Revisiting the fundamentals of phosphorus fractionation of sediments and soils. *J Soils Sediments*, 2011, 11:830-840.
- Dean L A. An attempted fractionation of the soil phosphorus. *Jour. Agr. Sci.*, 1938, 28: 234-246.
- Dorich, R. A., D. W. Nelson & L. E. Sommers. Estimating algal available phosphorus in suspended sediments by chemical extraction. *J. Environ. Qual.*, 1985, 14: 400-405.
- Fabre A, Qotbi A, Dauta A, Baldy V. Relation between algal available phosphate in the sediments of the River Garonne and chemically-determined phosphate fractions. *Hydrobiologia*, 1996, 335 : 43-48.
- Golterman H L. Differential extraction of sediment phosphates with NTA solution. *Hydrobiologia*, 1982, 92: 683-687.
- Golterman H L. The role of the ironhydroxide-phosphate-sulphide system in the phosphate exchange between sediments and overlying water. *Hydrobiologia*, 1995, 297 : 43-54.
- Golterman H L. Fractionation of sediment phosphate with chelating compounds: a simplification, and comparison with other methods. *Hydrobiologia*, 1996, 335: 87-95.
- Golterman H L, Paing J, Serrano L, Gomez E. Presence of and phosphate release from polyphosphates or phytate phosphate in lake sediments. *Hydrobiologia*, 1998, 364: 99-104.
- Hieltjes A H M, Lijklema L. Fractionation of inorganic phosphates in calcareous sediments. *J. environ. Qual.*, 1980, 9: 405-407.
- Hoyle C. Phosphorus Cycling in Lake Brienz, Switzerland. MSc thesis, EAWAG. ch, 14p, 2004.
- Hupfer M, Gächter R, Rüegger H. Polyphosphate in lake sediments: ^{31}P NMR spectroscopy as a tool for its identification. *Limnol. Oceanogr.*, 1995, 40(3): 610-617.
- Jaisi D P and Blake R E. Tracing sources and cycling of phosphorus in Peru Margin sediments using oxygen isotopes in authigenic and detrital phosphates. *Geochim. Cosmochim. Acta*, 2010, 74: 3199-3212.
- Jensen H S, Thamdrup B. Iron-bound phosphorus in marine sediments as measured by bicarbonate-dithionite extraction. *Hydrobiologia*, 1993, 253:47-59.
- Jensen H S, McGlathery K J, Marino R, Howarth R W. Forms and availability of sediment phosphorus in carbonate sand of Bermuda seagrass beds. *Limnol. Oceanogr.*, 1998, 43:799-810.

- Katsaounos C Z, Giokas D L, Leonardos J D and Karayannis M I. Speciation of phosphorus fractionation in river sediments by explanatory data analysis. *Water Res.*, 2007, 41: 406–418.
- Kerr J G, Burford M, Olley J and Udy J. The effects of drying on phosphorus sorption and speciation in subtropical river sediments. *Mar. Freshwater Res.*, 2010, 61, 928–935.
- Lukkari K, Hartikainen H and Leivuori M. Fractionation of sediment phosphorus revisited. I: Fractionation steps and their biogeochemical basis. *Limnol. Oceanogr.: Methods*, 2007, 5: 433–444.
- Monbet P, McKelvie I D. and Worsfold P J. Phosphorus speciation, burial and regeneration in coastal lagoon sediments of the Gippsland Lakes (Victoria, Australia). *Environ. Chem.*, 2007, 4: 334–346.
- Penn M R, Auer M T, Van Orman E L, Korienek J J. Phosphorus diagenesis in lake sediments: investigations using fractionation techniques. *Mar. Freshwater Res.*, 1995, 46:89-99.
- Penn M R, Auer M T. Seasonal variability in phosphorus speciation and deposition in a calcareous, eutrophic lake. *Mar. Geol.*, 1997, 139: 47-59.
- Pettersson K, Boström B and Jacobsen O. Phosphorus in sediments - speciation and analysis. *Hydrobiologia*, 1988, 170: 91-101.
- Psenner R, and Pucsko R. Phosphorus fractionation: advantages and limits of the method for the study of sediment P origins and interactions. *Arch. Hydrobiol. Beih. Ergebn. Limnol.*, 1988, 30:43-59.
- Psenner R, Bostrom B, Dinka M, Petterson K, Pucsko R, Sager M. Fractionation of phosphorus in suspended matter and sediment. *Arch. Hydrobiol. Beih.*, 1988, 30: 98-110.
- Ruban V, Brigault S, Demare D, Philippe A-M. An investigation of the origin and mobility of phosphorus in freshwater sediments from Bort-Les-Orgues Reservoir, France. *J. Environ. Monit.*, 1999a, 1, 403–407.
- Ruban V, López-Sánchez J F, Pardo P, Rauret G, Muntau H and Quevauviller P. Selection and evaluation of sequential extraction procedures for the determination of phosphorus forms in lake sediment. *J. Environ. Monit.*, 1999b, 1, 51–56.
- Ruban V, López-Sánchez J F, Pardo P, Rauret G, Muntau H and Quevauviller P. Harmonized protocol and certified reference material for the determination of extractable contents of phosphorus in freshwater sediments – A synthesis of recent works. *Fresenius J. Anal. Chem.*, 2001, 370: 224–228.
- Ruttenberg K C. Diagenesis and burial of phosphorus in marine sediments: Implications for the marine phosphorus budget. Ph.D. thesis, Yale Univ. 375 p. 1990.
- Ruttenberg K C. Development of a sequential extraction method for different forms of phosphorus in marine sediments. *Limnol. Oceanogr.*, 1992, 37: 1460-1482.
- Schindler D W, Kling H, Schmidt R V, Prokopowich J, Frost V E, Reid R A and Capel M. Eutrophication of Lake 227 by Addition of Phosphate and Nitrate: the Second, Third, and Fourth Years of Enrichment, 1970, 1971, and 1972. *J. Fish. Res. Board Can.*, 1973, 30(10): 1415-1440.
- Simon N S, Lynch D, Gallaher T N. Phosphorus Fractionation in Sediment Cores Collected In 2005 Before and After Onset of an *Aphanizomenon flos-aquae* Bloom in Upper Klamath Lake, OR, USA. *Water Air Soil Poll*, 2009, 204:139–153.
- Sharpley A. N. An innovative approach to estimate bioavailable phosphorus in agricultural runoff using iron-oxide impregnated paper. *J. Environ. Qual.*, 1993a, 22, 597-601.
- Sharpley A. N. Estimating phosphorus in agricultural runoff available to several algae using

- iron-oxide paper strips. *J. Environ. Qual.*, 1993b, 22, 678-680.
- Tiyapongpattana W, Pongsakul P, Shiowatana J and Nacapricha D. Sequential extraction of phosphorus in soil and sediment using a continuous-flow system. *Talanta*, 2004, 62(4): 765-771.
- Williams J D H, Syers a J K, Walter T W. Fractionation of soil inorganic phosphate by a modification of Chang and Jackson procedure. *Soil Sci. Soc. Amer. Proc.*, 1967, 31: 736-739.
- Williams J D H, Syers J K, Armstrong D E, Harris F. Fractionation of inorganic phosphate in calcareous lake sediments. *Soil Sci. Soc. Amer. Proc.*, 1971, 35: 250-255.
- Williams J D H, Jaquet J-M, Thomas R L. Forms of phosphorus in surficial sediments of Lake Erie. *J. Fish Res. Bd Can.*, 1976, 33: 413-429.
- Williams J D H, Shear H, Thomas R L. Availability to *Scenedesmus quadricauda* of different forms of phosphorus in sedimentary materials from the Great Lakes. *Limnol. oceanogr.*, 1980, 25 (1):1-11.
- Wang P, He M, Lin C, Men B, Liu R, Quan X and Yang Z. Phosphorus distribution in the estuarine sediments of the Daliao river, China. *Estuar. Coast. Shelf S.*, 2009, 84: 246-252.
- Yuan H Z, Shen J, Liu E F, Wang J J and Meng X H. Space Distribution characteristics and diversity analysis of phosphorus from overlying water and surface sediments in Taihu Lake. *Environ. Sci.*, 2010, 31: 954-960.
- Zhang J Z, Fischer C J and Ortner P B. Potential availability of sedimentary phosphorus to sediment resuspension in Florida Bay. *Global Biogeochem. Cy.*, 2004, 18:1-14.